

(d) *Directions.* The labeling of the product states, under the heading “Directions,” directions that conform to the directions established for each ingredient in the directions sections of this part. When the time intervals or age limitations for administration of the individual ingredients differ, the directions for the combination product may not exceed any maximum dosage limits established for the individual ingredients in the applicable OTC drug monograph.

§ 349.80 Professional labeling.

The labeling of any OTC ophthalmic demulcent drug product provided to health professionals (but not to the general public) may contain instructions for the use of these products in professional eye examinations (i.e. gonioscopy, electroretinography).

PART 350—ANTIPERSPIRANT DRUG PRODUCTS FOR OVER-THE-COUNTER HUMAN USE

Subpart A—General Provisions

- Sec.
350.1 Scope.
350.3 Definition.

Subpart B—Active Ingredients

- 350.10 Antiperspirant active ingredients.

Subpart C—Labeling

- 350.50 Labeling of antiperspirant drug products.

Subpart D—Guidelines for Effectiveness Testing

- 350.60 Guidelines for effectiveness testing of antiperspirant drug products.

AUTHORITY: 21 U.S.C. 321, 351, 352, 353, 355, 360, 371.

SOURCE: 68 FR 34291, June 9, 2003, unless otherwise noted.

Subpart A—General Provisions

§ 350.1 Scope.

(a) An over-the-counter antiperspirant drug product in a form suitable for topical administration is generally recognized as safe and effective and is not misbranded if it meets each condition in this part and each general

condition established in § 330.1 of this chapter.

(b) References in this part to regulatory sections of the Code of Federal Regulations are to chapter I of title 21 unless otherwise noted.

§ 350.3 Definition.

As used in this part:

Antiperspirant. A drug product applied topically that reduces the production of perspiration (sweat) at that site.

Subpart B—Active Ingredients

§ 350.10 Antiperspirant active ingredients.

The active ingredient of the product consists of any of the following within the established concentration and dosage formulation. Where applicable, the ingredient must meet the aluminum to chloride, aluminum to zirconium, and aluminum plus zirconium to chloride atomic ratios described in the U.S. Pharmacopeia-National Formulary. The concentration of ingredients in paragraphs (b) through (j) of this section is calculated on an anhydrous basis, omitting from the calculation any buffer component present in the compound, in an aerosol or nonaerosol dosage form. The concentration of ingredients in paragraphs (k) through (r) of this section is calculated on an anhydrous basis, omitting from the calculation any buffer component present in the compound, in a nonaerosol dosage form. The labeled declaration of the percentage of the active ingredient should exclude any water, buffer components, or propellant.

(a) Aluminum chloride up to 15 percent, calculated on the hexahydrate form, in an aqueous solution nonaerosol dosage form.

(b) Aluminum chlorohydrate up to 25 percent.

(c) Aluminum chlorohydrate polyethylene glycol up to 25 percent.

(d) Aluminum chlorohydrate propylene glycol up to 25 percent.

(e) Aluminum dichlorohydrate up to 25 percent.

(f) Aluminum dichlorohydrate polyethylene glycol up to 25 percent.

(g) Aluminum dichlorohydrate propylene glycol up to 25 percent.